## This Page Is Inserted by IFW Operations and is not a part of the Official Record

#### **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


### IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.


# IgG WESTERN BLOT WITH 5 DIFFERENT STRAINS AS ANTIGEN IgG AND IgM RESPONSE IN STAGE III

NEUROBORRELIOSIS, STAGE II (IgM) (IgG)

Neuroborreliose, Stadium II (IgM) (IgG)



### IgM WESTERN BLOT WHEN THERE IS ERYTHEMA MIGRANS ANTIGEN: B. BURGDORFERI/PKo STRAIN



PATIENT A: 1 FIRST TEST


2,3 MONITORING PROGRESS AFTER 1 AND 3 WEEKS

PATIENT B: 1 FIRST TEST

2-5 MONITORING PROGRESS AFTER 2, 9, 10 AND 11 WEEKS


FIG. 2a


# IgM WESTERN BLOT WHEN THERE ARE LATE MANIFESTATIONS ANTIGEN: B. BURGDORFERI/PKo STRAIN



1, 2, 3 ACRODERMATITIS 4, 5 ARTHRITIS

FIG. 2b










FIG. 5



MONOCLONAL ANTIBODIES AGAINST B. BURGDORFERI



, MONOCLONAL ANTIBODIES AGAINST B. BURGDORFERI

